

## Math 230H, Extra Credit Problems

These are hard and interesting problems. It might improve your score, but should be used for fun. Only the first solution will be graded. The solutions should be presents orally before November 18.

1. Given  $n$  vectors  $\mathbf{v}_1, \dots, \mathbf{v}_n$ , denote by  $A$  the sum  $n$  numbers  $\|\mathbf{v}_i\|^2$  and by  $B$  the sum of  $n(n-1)/2$  numbers  $\|\mathbf{v}_i - \mathbf{v}_j\|^2$  for  $1 \leq i < j \leq n$ .

Show that

$$nA \geq B.$$

2. Two sets  $\Gamma$  and  $\Delta$  in the Euclidean space are called *equidistant* if the distance function from  $\Gamma$  is constant on  $\Delta$  and the distance function from  $\Delta$  is constant on  $\Gamma$ . (For example, two concentric circles in one plane are equidistant.)

Describe all the pairs of equidistant circles in the Euclidean space.

- 3a. (Solved) Given two vectors  $\mathbf{x}_0$  and  $\mathbf{v}$ , describe geometrically the set of solutions of the vector equation

$$\mathbf{x} \times (\mathbf{x} \times \mathbf{v}) = \mathbf{x}_0 \times (\mathbf{x}_0 \times \mathbf{v})$$

with unknown vector  $\mathbf{x}$ .

- 3b. (Solved) Given two vectors  $\mathbf{v}$  and  $\mathbf{w}$ , describe geometrically the set of solutions of the vector equation

$$(\mathbf{x} \times \mathbf{v}) \times (\mathbf{x} \times \mathbf{w}) = \mathbf{0}$$

with unknown vector  $\mathbf{x}$ .

- 3c. (Solved) Given two vectors  $\mathbf{v}$  and  $\mathbf{w}$ , describe geometrically the set of solutions of the vector equation

$$\mathbf{x} \times \mathbf{v} = \|\mathbf{x}\|^2 \mathbf{w}$$

with unknown vector  $\mathbf{x}$ .

4. (Solved) Given two points  $P$  and  $Q$  in the space consider the set of points  $X$  such that the distance from  $X$  to  $P$  is twice larger than the distance from  $X$  to

Q. Show that the set is formed by a sphere, find its radius and center in terms of  $P$  and  $Q$ .

5. Show that equidistant set from two skew lines is a hyperbolic paraboloid.

6. (Solved) Show that there is no plane which is *tangent* to the curve

$$\mathbf{f}(t) = (t, t^2, t^3)$$

at two distinct points.

(A plane is called *tangent* to a curve  $\mathbf{f}(t)$  at point  $\mathbf{f}(t_0)$  if it contains the tangent line at  $\mathbf{f}(t_0)$ .)

7a. Let  $\mathbf{f}(t)$  be a smooth curve in the plane. Assume its curvature  $\kappa(t)$  is increasing in  $t$ . Show that the curve has *no self-intersections*; that is, if  $t_0 \neq t_1$  then  $\mathbf{f}(t_0) \neq \mathbf{f}(t_1)$ .

7b. Assume  $\mathbf{f}(t)$  is a closed smooth curve in the unit ball with a unit-speed parametrization. Show that the average of curvature of  $\mathbf{f}(t)$  is at least 1.

7c. Assume  $\mathbf{f}(t)$  is a periodic closed space curve a unit-speed parametrization. Show that the integral of curvature is at least  $2\pi$ .

8. Assume  $f$  is a convex function of two variables. Let  $\mathbf{x}(t) = (x_1(t), x_2(t))$  and  $\mathbf{y}(t) = (y_1(t), y_2(t))$  be two plane curves such that

$$\mathbf{x}'(t) = \nabla f(\mathbf{x}(t)) \quad \text{and} \quad \mathbf{y}'(t) = \nabla f(\mathbf{y}(t))$$

for any  $t$ . Show that the function

$$\ell(t) = \|\mathbf{x}(t) - \mathbf{y}(t)\|$$

is nondecreasing.

(A function  $f(x, y)$  is called *convex* if its epigraph  $z \geq f(x, y)$  is a convex set.)

9. Let  $f$  be a smooth function of two variables. Assume that  $\mathbf{x}(t)$  is a smooth closed plane curve such that

$$D_{\mathbf{x}'(t)}^2 f(\mathbf{x}(t)) = 0$$


for any  $t$  and

$$\frac{\partial^2 f}{\partial x^2} \frac{\partial^2 f}{\partial y^2} - \left(\frac{\partial^2 f}{\partial x \partial y}\right)^2 < 0$$

at any point  $\mathbf{x}(t)$ .

Show that the curve  $\mathbf{x}(t)$  can not be star-shaped.

(A closed plane curve is called *star-shaped* if every ray from some fixed point  $P$  intersects the curve at a single point.)



10. Let  $(x_0, y_0)$  be a minimum point of smooth function  $f(x, y)$  with the constraint  $g(x, y) \leq c$ . Assume  $g(x_0, y_0) = c$  and  $\nabla g(x_0, y_0) \neq \mathbf{0}$ . Show that  $\nabla f(x_0, y_0) = \lambda \nabla g(x_0, y_0)$  for some  $\lambda \leq 0$ .

11. Let  $F$  be a figure in the upper half-plane; denote as  $(x_0, y_0)$  its center of mass and as  $A$  its the area. Show that

$$2\pi A y_0$$

is the volume of the body of revolution obtained by rotating  $F$  around  $x$ -axis.

~~12.~~ (Solved) Consider the vector field  $\mathbf{f}(x, y, z) = (0, x, 1)$ . Show that any two points in the space can be connected by a curve  $\mathbf{r}(t) = (x(t), y(t), z(t))$  such that

$$\mathbf{r}'(t) \perp \mathbf{f}(x(t), y(t), z(t))$$

for any  $t$ .

13. Assume  $K$  a body is bounded by two figures in parallel planes and the ruled surface containing the curves bounding both figures. Let  $h$  be the height of  $K$ ,  $A_-$  be the area of the lower base of  $K$ ,  $A_0$  be the area of the middle section of  $K$  and  $A_+$  be the area of the upper base of  $K$ . Show that

$$\frac{h}{6}(A_- + 4A_0 + A_+)$$

is the volume of  $K$ .